

A COMBINATORIAL APPROACH TO ALEXANDER–HIRSCHOWITZ’S THEOREM BASED ON TORIC DEGENERATIONS

SILVIA BRANNETTI

ABSTRACT. We present an alternative proof of the Alexander–Hirschowitz’s Theorem in dimension 3 using degenerations of toric varieties.

1. INTRODUCTION

Secant varieties are classical objects of study in algebraic geometry: given a closed variety X in some projective space $\mathbb{P}^n$, and given a natural number k , the k -th secant variety of X is the Zariski closure of the union of all subspaces of $\mathbb{P}^n$ that are spanned by $k + 1$ independent points on X . Not enough is known about these varieties, starting from their dimensions. Cases of interest are for instance the secant varieties of Segre embeddings of products of projective spaces, Plücker embeddings of Grassmannians, and Veronese embeddings of projective spaces. The Alexander–Hirschowitz’s Theorem (2.3) provides a complete answer to the case of all Veronese embeddings of $\mathbb{P}^n$. We recall that the problem of the secant varieties is equivalent to another classical one in Algebraic Geometry, Polynomial Interpolation (see Remark 2.4), and in this setting Alexander–Hirschowitz’s Theorem solves exhaustively the problem on interpolation with double points in any dimension; its original proof is due to Alexander and Hirschowitz in the nineties, and in 2002 it has been simplified by Chandler [Ch02]. More recently the proof has been improved by Brambilla and Ottaviani in [BO07]. Moreover a proof of this theorem has been given in dimension 2 by Draisma ([JD06]) using a new approach based on tropical geometry, and again in dimension 2 a different proof of the theorem is by Ciliberto, Miranda, Dumitrescu ([CM06]) involving degenerations of toric varieties, that allow to translate the classical problem into an easy combinatorial one. Indeed tropical geometry is closely related to degenerations of toric varieties, so that the two methods are in fact very much connected. For related concepts see ([SS05]). In this paper we want to prove Alexander–Hirschowitz’s Theorem in dimension 3 applying toric degenerations, in line with the approach of [CM06], and exploiting the combinatorial nature of toric varieties.

ACKNOWLEDGEMENTS

I wish to thank Prof. Ciro Ciliberto for several useful discussions, precious suggestions and patient encouragements. I also thank the referee of the paper for his very detailed report and his advices.

2. SECANT VARIETIES AND THE ALEXANDER–HIRSCHOWITZ’S THEOREM

Definition 2.1. Let $X \subset \mathbb{P}^N$ be an irreducible, non-degenerate projective variety of dimension n and let k be a positive integer. Take $k + 1$ independent points $p_0, \dots, p_k$ of X . The span $\langle p_0, \dots, p_k \rangle$ is a subspace of $\mathbb{P}^N$ of dimension k which is called a $(k + 1)$ -secant $\mathbb{P}^k$ of X . By $\text{Sec}_k(X)$ we denote the closure of the union of all $(k + 1)$ -secant $\mathbb{P}^k$ ’s of X . This is an irreducible algebraic variety which is called the k -th secant variety of X .

The study of secant varieties in particular concerns their dimension, which in most cases is unknown. It is easy to see that there is a natural upper bound on the dimension:

$$(2.1) \quad \dim(\text{Sec}_k(X)) \leq \min\{(n + 1)(k + 1) - 1, N\}.$$

The right hand side of (2.1) is the *expected dimension* of $\text{Sec}_k(X)$.

Definition 2.2. X is said to be k -defective if the strict inequality holds in (2.1), and its k -defect is $\delta_k = \min\{(n + 1)(k + 1) - 1, N\} - \dim(\text{Sec}_k(X))$, otherwise it is said to be not k -defective.

Our aim in this paper is to prove Alexander–Hirschowitz’s Theorem in dimension 3, so we are going to recall its terms. Let $V_{n,d}$ be the Veronese variety in $\mathbb{P}^{\binom{n+d}{d}-1}$:

Theorem 2.3. (Alexander–Hirschowitz) *The Veronese variety $V_{n,d}$ is always not k -defective, except in the cases:*

n	any	2	3	4	4
d	2	4	4	4	3
k	$2, \dots, n$	5	9	14	7.

Terracini’s lemma (see [Ci01]) allows us to reduce the problem of computing the dimension of $\text{Sec}_k(X)$ to the study of the dimension of the span of the linear spaces T_{X,p_i} , $i = 0, \dots, k$ which are the tangent spaces to X at $p_0, \dots, p_k$, for $p_0, \dots, p_k$ general points on X . Let us observe that if the T_{X,p_i} with $i = 0, \dots, k$ are independent in $\mathbb{P}^N$, then their span has maximal dimension, equal to $(n+1)(k+1) - 1$, therefore X is not k -defective. Since our aim is to show Alexander–Hirschowitz’s Theorem in dimension 3, we are interested in studying the defectivity of the secant variety of the Veronese threefold $V_{3,d}$, which is isomorphic to $\mathbb{P}^3$ via the Veronese embedding. We want to show that $V_{3,d}$ is not k -defective, for any $k \in \mathbb{N}$ and $d \geq 5$. The cases $d \leq 4$ are well known. In general, given a variety X as in Definition 2.1, let $\mathcal{D}$ be a degeneration of X to a union of varieties $X_1, \dots, X_m$; fix $p_0, \dots, p_k$ general points on X . If we let the $k+1$ points degenerate in such a way that each point goes to a general point of some component of the central fiber, we can consider the tangent spaces to the central fiber at the limit points. We denote by $T_{X_i,p}$ the tangent space to the component X_i of the central fiber at p , where p is a limit point belonging to X_i . If the tangent spaces to the central fiber at the limit points are independent, so are the tangent spaces T_{X,p_i} , $i = 0, \dots, k$, hence X is not k -defective.

Remark 2.4. We explain briefly the connections of the problem of secant varieties with polynomial interpolation. The latter can be stated as follows: given a set of points $p_0, \dots, p_k$ in $\mathbb{P}^n$, and assigned relative multiplicities $m_0, \dots, m_k \in \mathbb{N}$, we study the dimension of the linear system $\mathcal{L}_{n,d}(\sum_{i=0}^k m_i p_i)$ of the hypersurfaces of degree d in $\mathbb{P}^n$ having at least multiplicity m_i at p_i for each $i = 0, \dots, k$. The system is said to be *special* if it doesn’t have the expected dimension, otherwise it is said to be *non-special*. The study of interpolation problem consists in classifying the special systems. This has been completely done for $n = 1$, $n = 2$ and for any n when the multiplicities are $m_i = 2 \forall i$. The latter case is examined in Alexander–Hirschowitz’s Theorem, as noted above. By Terracini’s lemma we can say that $\text{Sec}_k(V_{n,d})$ has the expected dimension if and only if the system $\mathcal{L}_{n,d}(2^{k+1})$ is non-special, indeed the Veronese morphism $v_{n,d}$ allows us to embed $\mathbb{P}^n$ in $\mathbb{P}^{\binom{n+d}{d}-1}$, and to translate the system of the hypersurfaces of degree d with $k+1$ double points, to the system of hyperplanes in $\mathbb{P}^{\binom{n+d}{d}-1}$ which are tangent to $V_{n,d}$ at $k+1$ fixed general points. In [CM06] the authors prove Alexander–Hirschowitz’s Theorem in dimension 2, using suitable degenerations of the Veronese surface $V_{2,d}$, and reducing the problem to an easy combinatorial one. Even if one could use the same technique in dimension three, we will give a similar approach that emphasizes the perspective of toric varieties as described below.

2.1. Degenerations of toric varieties. In this paper we will use degenerations of *toric varieties*. Let us briefly recall how a degeneration of a toric variety can be described. The interested reader is referred to [Hu01] and [CM06] for details. A rational polytope $\mathcal{P}$ in $\mathbb{R}^n$ defines a projective toric variety $X_{\mathcal{P}}$ endowed with an ample line bundle; a subdivision Γ of $\mathcal{P}$ is a partition of $\mathcal{P}$ into smaller polytopes, i.e. there exist polytopes $\mathcal{P}_1, \dots, \mathcal{P}_l$ of the same dimension as $\mathcal{P}$, such that $\mathcal{P}_i \cap \mathcal{P}_j$ is a face of both (it can be the empty face), and $\bigcup_{i=1}^l \mathcal{P}_i = \mathcal{P}$. Then Γ is said to be a *regular subdivision* of $\mathcal{P}$ if there exists an integral function F defined on $\mathcal{P}$, which is piecewise linear on the subpolytopes of Γ and strictly convex on Γ , i.e. for any points p and q in different subpolytopes of Γ , $F(tp + (1-t)q) < tF(p) + (1-t)F(q)$ for all $t \in [0, 1]$. If Γ is regular, then there exists a degeneration of $X_{\mathcal{P}}$ such that the central fiber is the divisor X_0 whose components are the toric varieties defined by the subpolytopes of Γ , whereas the total space of the degeneration is the toric variety associated to the unbounded polytope $\tilde{\mathcal{P}} = \{(x, z) \in \mathcal{P} \times \mathbb{R} : z \geq F(x)\}$. In this construction $X_{\mathcal{P}}$ is isomorphic to the general fiber, thus it degenerates to X_0 . If X_i is a component of the central fiber endowed with a very ample line bundle, then the corresponding subpolytope $\mathcal{P}_i$ provides an embedding of X_i in the projective space $\mathbb{P}^{l_i-1}$, where l_i is the number of integral points in $\mathcal{P}_i$. Note that if two subpolytopes $\mathcal{P}_i, \mathcal{P}_j$ in a subdivision Γ of $\mathcal{P}$ are disjoint, then the varieties they define are embedded in independent linear subspaces of $\mathbb{P}^{M-1}$, where M is the number of integral points of $\mathcal{P}$.

Proposition 2.5. *Let $X \subset \mathbb{P}^N$ be an irreducible, toric projective variety of dimension n , let $\mathcal{D}$ be a toric degeneration of X to $X_0 = X_1 \cup \dots \cup X_m$, and let $\mathcal{P}_1, \dots, \mathcal{P}_m$ be the corresponding polytopes. If*

- (i) *there exist indices $j_1, \dots, j_h \in \{1, \dots, m\}$ such that X_{j_i} is not k_{j_i} -defective,*
- (ii) *$\mathcal{P}_{j_i} \cap \mathcal{P}_{j_l} = \emptyset$ for any $i \neq l$, with $i, l = 1, \dots, h$,*

then, setting $k+1 = \sum_{i=1}^h (k_{j_i} + 1)$, X is not k -defective.

Proof. We can assume $j_1 = 1, \dots, j_h = h$, so X_i is not k_i -defective, for any $i = 1, \dots, h$; fix p_j^i , $j = 0, \dots, k_i$ general points on X_i . Then $\dim(\langle \bigcup_{j=0, \dots, k_i} T_{X_i, p_j^i} \rangle) = (n+1)(k_i+1) - 1$. Since the polytopes $\mathcal{P}_1, \dots, \mathcal{P}_h$ are disjoint, then we can conclude that

$$\dim(\langle \bigcup_{\substack{j=0, \dots, k_i \\ i=1, \dots, h}} T_{X_i, p_j^i} \rangle) = (n+1)(k+1) - 1.$$

We end the proof by observing that the points p_j^i are limits of general points on X . □

Now let X be the Veronese threefold $V_{3,d}$ in $\mathbb{P}^{N_d}$, where $N_d = \binom{d+3}{3} - 1$. We want to prove Alexander-Hirschowitz's Theorem for any $d \geq 5$ and any $k \in \mathbb{N}$. Let $n_d + 1 = \left\lfloor \frac{(d+1)(d+2)(d+3)}{24} \right\rfloor$; we have the following results, whose easy proofs can be left to the reader:

Lemma 2.6. *The expected dimension of $\text{Sec}_{n_d}(V_{3,d})$ is*

$$(2.2) \quad \begin{cases} N_d & \text{if } d \text{ is odd or } d = 6 + 8k, \text{ for } k \geq 0 \\ N_d - 1 & \text{if } d = 8k, \text{ for } k \geq 1 \\ N_d - 2 & \text{if } d = 10 + 8k, \text{ for } k \geq 0 \\ N_d - 3 & \text{if } d = 12 + 8k, \text{ for } k \geq 0. \end{cases}$$

Lemma 2.7. *If $V_{3,d}$ is not n_d -defective for any $d \geq 5$, then it is not k -defective for any $k < n_d$.*

Remark 2.8. Suppose we have proved that $V_{3,d}$ is not n_d -defective. If $k > n_d$, in order to see that $V_{3,d}$ is not k -defective we have to prove that the dimension of the span of the tangent spaces at $k+1$ general points is maximal. By Lemma 2.6 this is obvious when either d is odd, or when $d \equiv 6 \pmod{8}$; in the other cases we have that $\text{Sec}_{n_d}(V_{3,d})$ has codimension 1, 2 or 3 depending on $d \equiv 0, 2$ or $4 \pmod{8}$. If we impose one further general point in such a way that its tangent space and the tangent spaces to the points that we have already imposed span the whole ambient space $\mathbb{P}^{N_d}$, the dimension of $\text{Sec}_{n_d}(V_{3,d})$ is maximal.

The degeneration of the Veronese threefold we will deal with, is described in the following:

Lemma 2.9. *The Veronese threefold $V_{3,d}$ degenerates to $\binom{d}{3}$ copies of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, plus $\frac{d(d+1)}{2}$ copies of $\mathbb{P}^3$, and $\frac{d(d-1)}{2}$ copies of the blow up of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ at some point.*






FIGURE 1. Δ_d

Proof. Let Δ_d be the polytope defining $V_{3,d}$ as a toric variety; it is a tetrahedron of side length d , as in Figure 1. We start cutting Δ_d horizontally, so we get a Δ_{d-1} leaning on one layer of height 1, as in Figure 2. We denote by S_k^1 a layer of height 1 and base of side length k . Iterating the cut in Δ_{d-1} we get a subdivision of Δ_d in $(d-1)$ layers of height 1 plus a tetrahedron on the top. Then we further subdivide each layer into a certain number of polytopes, i.e. inside the layer S_k^1 there will be $\frac{(k-1)(k-2)}{2}$ cubes, k tetrahedra Δ_1 , and $k-1$ blocks Σ as the one represented in Figure 3. The polytope Σ can be interpreted as obtained by removing a tetrahedron from a cube; this corresponds to the blow up of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ at one point, which we denote $(\widehat{\mathbb{P}^1})^3$. So far we obtained a subdivision Γ of Δ_d into cubes, tetrahedra Δ_1 and blocks Σ . Let C_{ijk} be the cube having vertices $(i, j, k), (i+1, j, k), (i, j+1, k), (i, j, k+1), (i+1, j+1, k), (i+1, j, k+1), (i, j+1, k+1), (i+1, j+1, k+1)$, with $i, j, k \in \mathbb{N}$. Our piecewise linear function F is defined as follows: on the cube C_{ijk} in the partition of Δ_d , the function is given by the linear form

$$F|_{C_{ijk}}(x_1, x_2, x_3) = (1+2i)x_1 + (1+2j)x_2 + (1+2k)x_3 - (i+j+k+i^2+j^2+k^2).$$


FIGURE 2. Section of Δ_d into Δ_{d-1} plus one layer of height 1.FIGURE 3. The block Σ .

An easy calculation shows that the function is strictly convex on our subdivision Γ . Hence the subdivision Γ is regular, and it yields a degeneration of $V_{3,d}$ to $\binom{d}{3}$ copies of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, corresponding to the cubes, plus $\frac{d(d+1)}{2}$ copies of $\mathbb{P}^3$, corresponding to the tetrahedra, and $\frac{d(d-1)}{2}$ copies of the blow up of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ at some point, corresponding to the blocks Σ . □

In the following we are going to prove Alexander–Hirschowitz’s Theorem treating in separated sections the cases d odd and d even. We start with the following Lemmata, which will be useful throughout.

Lemma 2.10. *Let C be the unitary cube in $\mathbb{R}^3$, and let T be one of the 8 tetrahedra Δ_1 containing 3 edges of C meeting at one vertex. Then the toric variety it defines is a tangent $\mathbb{P}^3$ to $(\mathbb{P}^1)^3$ at one coordinate point of $\mathbb{P}^7$.*

Proof. Note that T is a tetrahedron like the one in Figure 4. We can assume that C is the cube with the origin as a vertex, and that T is the tetrahedron of vertices $\{(0, 0, 0), (1, 0, 0), (0, 1, 0), (0, 0, 1)\}$. Then, an immediate computation shows that T is the tangent space to the Segre variety $(\mathbb{P}^1)^3$ defined by C , at $[1 : 0 : 0 : 0 : 0 : 0 : 0 : 0]$, one of the coordinate points of the $\mathbb{P}^7$ spanned by $(\mathbb{P}^1)^3$. □

FIGURE 4. One of the 8 tetrahedra Δ_1 containing 3 edges of C meeting at one vertex.

Lemma 2.11. *The following facts hold:*

- (i) $\mathbb{P}^3$ is not 0-defective.
- (ii) The Segre threefold $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ is not 1-defective.

Proof. Assertion (i) is obvious by Definition 2.2, since $\text{Sec}_0(\mathbb{P}^3) = \mathbb{P}^3$. In order to prove (ii), we recall that the Segre embedding of $(\mathbb{P}^1)^3$ in $\mathbb{P}^7$ is represented by a cube as a toric variety. If we look at Figure 5, we see that the




FIGURE 5. The subdivision of the cube.

two tetrahedra pointed out are of the type introduced in Lemma 2.10, i.e. they correspond to tangent $\mathbb{P}^3$ ’s to $(\mathbb{P}^1)^3$.

Since they are disjoint, the relative tangent $\mathbb{P}^3$ ’s are skew in the ambient space $\mathbb{P}^7$, so they span a linear space of maximal dimension $4 \cdot 2 - 1 = 7$. If instead of the tangent $\mathbb{P}^3$ ’s defined by the tetrahedra in the picture we take general ones, by semicontinuity we get that $\dim(\text{Sec}_1(\mathbb{P}^1)^3)$ is maximal, hence $(\mathbb{P}^1)^3$ is not 1–defective. $\square$

3. CASE d ODD

Let d be odd, and let $\mathcal{D}$ be the degeneration of $V_{3,d}$ as in Lemma 2.9; by the same Lemma we have that each layer S_k^1 degenerates to $\frac{(k-1)(k-2)}{2}$ copies of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, k copies of $\mathbb{P}^3$ and $k-1$ copies of $(\mathbb{P}^1)^3$. We recall that a nonsingular polytope Δ describes a closed embedding of the toric variety $X(\Delta)$, i.e. it determines a very ample line bundle on $X(\Delta)$ [Fu93]. Then S_k^1 corresponds to $\mathbb{P}^3$ blown up at a point p , embedded in $\mathbb{P}^{M_k-1}$, where $M_k = \frac{(k+2)(k+1)}{2} + \frac{(k+1)k}{2}$, via the proper transform of the linear system of surfaces of degree k with a point of multiplicity $k-1$ at p . If we denote by π the blow up, the line bundle which embeds $\widetilde{\mathbb{P}^3}$ in $\mathbb{P}^{M_k-1}$ is $\pi^*(\mathcal{O}_{\mathbb{P}^3}(k)) \otimes \mathcal{O}_{\widetilde{\mathbb{P}^3}}(-(k-1)E)$, where E is the exceptional divisor of the blow up. We have the following:

Lemma 3.1. *Let k be odd; then $\widetilde{\mathbb{P}^3}$, embedded in $\mathbb{P}^{M_k-1}$, is not $\left(\frac{(k+1)^2}{4} - 1\right)$ –defective.*

Proof. Consider the subdivision Γ_k of S_k^1 in cubes, Σ ’s and Δ_1 ’s. We have seen that it is regular; the unitary cube corresponds to the toric variety $(\mathbb{P}^1)^3$ which is not 1–defective, whereas the tetrahedron Δ_1 defines $\mathbb{P}^3$ as a not 0–defective variety (see Lemma 2.11). If we take disjoint polytopes as in Figure 6, we can apply Proposition 2.5. Indeed, in the layer S_k^1 we can point out $\frac{k^2-1}{8}$ disjoint cubes, and $\frac{k+1}{2}$ disjoint tetrahedra Δ_1 . Thus applying Proposition 2.5 we get that $\widetilde{\mathbb{P}^3}$ in its embedding in $\mathbb{P}^{M_k-1}$ is not $\left(2\frac{k^2-1}{8} + \frac{k+1}{2} - 1\right)$ –defective.




FIGURE 6. Projection of S_k^1

We are now ready to state:

Lemma 3.2. *Let d be odd, then $V_{3,d}$ is not n_d –defective.*

Proof. Consider the subdivision of Δ_d into $d-1$ layers S_k^1 plus one Δ_1 . This subdivision is regular by an argument analogous to the one used in Lemma 2.9. By the previous Lemma we know that S_k^1 for k odd is not $\left(\frac{(k+1)^2}{4} - 1\right)$ –defective. Let us take all the layers S_k^1 for $k \geq 3$ odd, and the tetrahedron Δ_1 on the top: they are disjoint, therefore applying Proposition 2.5, by induction on d we get that $V_{3,d}$ is not n_d –defective. Indeed, let $d = 5$: it is easy to see that $V_{3,5}$ is not 13–defective, since S_5^1 is not 8–defective, S_3^1 is not 3–defective and Δ_1 is not 0–defective. Suppose now that we have proved that $V_{3,d-2}$ is not n_{d-2} –defective. We obtain the tetrahedron Δ_d adding to Δ_{d-2} the layers S_{d-1}^1 and S_d^1 . So we just have to verify that $n_d + 1 = n_{d-2} + 1 + \frac{(d+1)^2}{4}$, which is an easy computation. $\square$

Remark 3.3. To see that $V_{3,d}$ for d odd is not k –defective for any $k \in \mathbb{N}$, we just have to apply Lemma 2.7 and Remark 2.8, since we have shown that $V_{3,d}$ is not n_d –defective.

4. CASE d EVEN

When d is even we cannot apply the same argument as for d odd because the situation is more complicated; by Lemma 2.6 we have to treat the cases $d = 6, 8, 10, 12$. Henceforth we will often regard the polytopes and the toric varieties they determine as the same objects.

Lemma 4.1. *$V_{3,6}$ is not 20–defective.*

Proof. Looking at Figure 7 one sees that among the 21 tetrahedra that we colored, 5 are not of the type introduced in Lemma 2.10; we denote them by $\Sigma_1, \dots, \Sigma_5$. They correspond to $\mathbb{P}^3$ as toric varieties, but they are not necessarily tangent $\mathbb{P}^3$ ’s. It is clear that the 21 $\mathbb{P}^3$ ’s are independent, since their polytopes are disjoint, so they span a projective space of maximal dimension, 83, that we denote by $\mathbb{P}^M$. The four vertices of each tetrahedron correspond to independent points in the ambient space, so we can think of $\mathbb{P}^M$ as the span of the 4–tuples of vertices of the 21 tetrahedra. Let us observe that a projectivity of $\mathbb{P}^M$ can be represented by a (84×84) matrix up to multiplication by scalars. Fix $i = 1, \dots, 5$, for example $i = 1$, and consider Σ_1 ; we have to distinguish two cases.




FIGURE 7. Δ_6 : it is decomposed in layers of height 1 in which we have pointed out pairwise disjoint tetrahedra.

Case I: the tetrahedron Σ_1 lies inside a cube. We define φ to be the projectivity associated to a diagonal matrix M_φ , whose diagonal is the string $(1, 1, 1, 1, t, \dots, t)$, where the first 4 columns of M_φ correspond to the 4 points generating Σ_1 , and the remaining columns, 4 by 4 correspond to the ordered 4-tuples of points spanning the other $\mathbb{P}^3$'s. As we know Σ_1 lies inside a cube, which corresponds to a $(\mathbb{P}^1)^3$; let H_1 be a general $\mathbb{P}^3$ tangent to the $(\mathbb{P}^1)^3$ where Σ_1 sits, at some general point. Since $(\mathbb{P}^1)^3$ spans a $\mathbb{P}^7$, we can choose points $p_1, \dots, p_4 \in H_1$ such that $\langle p_1, \dots, p_4 \rangle = H_1$, and such that $p_i = [a_{i0} : \dots : a_{i7} : 0 : \dots : 0]$ for any $i = 1, \dots, 4$. Applying now φ to H_1 , we get:

$$(4.1) \quad \varphi \begin{pmatrix} a_{10} & a_{20} & a_{30} & a_{40} \\ \cdot & \cdot & \cdot & \cdot \\ a_{17} & a_{27} & a_{37} & a_{47} \\ 0 & 0 & 0 & 0 \\ \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} a_{10} & a_{20} & a_{30} & a_{40} \\ \cdot & \cdot & \cdot & \cdot \\ a_{13} & a_{23} & a_{33} & a_{43} \\ ta_{14} & ta_{24} & ta_{34} & ta_{44} \\ \cdot & \cdot & \cdot & \cdot \\ ta_{17} & ta_{27} & ta_{37} & ta_{47} \\ 0 & 0 & 0 & 0 \\ \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Taking the limit as $t \rightarrow 0$ we obtain that the limit image of H_1 via φ is the space $\mathcal{H}_1$ defined by the following matrix

$$\begin{pmatrix} a_{10} & a_{20} & a_{30} & a_{40} \\ \vdots & \vdots & \vdots & \vdots \\ a_{13} & a_{23} & a_{33} & a_{43} \\ 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

We note that $\mathcal{H}_1$ is the image of H_1 via the projection $\pi : \mathbb{P}^7 \dashrightarrow \mathbb{P}^3$, such that $\pi(x_0 : \dots : x_7) \mapsto (x_0 : \dots : x_3)$, which is the projection from the $\mathbb{P}^3$ spanning with Σ_1 the $\mathbb{P}^7$ where $(\mathbb{P}^1)^3$ sits, i.e., the $\mathbb{P}^3$ generated by the 4 vertices of the cube not belonging to Σ_1 . An easy calculation shows that the general $\mathbb{P}^3$ tangent to $(\mathbb{P}^1)^3$, in particular H_1 , doesn’t intersect the center of the projection. Hence $\mathcal{H}_1$ is still a $\mathbb{P}^3$, and it is precisely Σ_1 .

Case II: The tetrahedron Σ_1 lies inside a block Σ as in Figure 3, so the $\mathbb{P}^3$ it defines sits in a $\mathbb{P}^6$, since the toric variety defined by Σ is a $(\mathbb{P}^1)^3$ blown up at a point. Let ψ be the projectivity of $\mathbb{P}^M$ defined by a diagonal matrix M_ψ , whose diagonal is the string $(1, 1, 1, 1, t, \dots, t)$, as well as in Case I, where the first four columns correspond to the four points generating Σ_1 . Let H_1 be a general $\mathbb{P}^3$ tangent to the $(\mathbb{P}^1)^3$ where the $\mathbb{P}^3$ corresponding to Σ_1 sits; choose $p_1, \dots, p_4 \in H_1$ such that $\langle p_1, \dots, p_4 \rangle = H_1$. The space H_1 is tangent to $(\mathbb{P}^1)^3$, which spans a $\mathbb{P}^6$, hence we can take the points $p_1, \dots, p_4$ such that $p_i = [a_{i0} : \dots : a_{i6} : 0 : \dots : 0]$ for any $i = 1, \dots, 4$. In this setting we use the argument of Case I, and obtain that the limit of H_1 via ψ is its image via the projection $\pi : \mathbb{P}^6 \dashrightarrow \mathbb{P}^3$, such that $\pi(x_0 : \dots : x_6) \mapsto (x_0 : \dots : x_3)$, which is the projection from the $\mathbb{P}^2$ spanning with Σ_1 the $\mathbb{P}^6$ where the blow up of $(\mathbb{P}^1)^3$ sits, i.e. the $\mathbb{P}^2$ generated by the 3 vertices of the block Σ not belonging to Σ_1 . An easy calculation shows that the general tangent $\mathbb{P}^3$ doesn’t intersect the center of projection, hence the limit of H_1 is still a $\mathbb{P}^3$, and it is exactly the one determined by Σ_1 .

Let us denote by Σ_5 the tetrahedron in Figure 7 that we see in the base layer S_6^1 , lying in a block Σ . The other tetrahedra, $\Sigma_1, \dots, \Sigma_4$, lie inside unitary cubes. We apply the construction of Case I to $\Sigma_1, \dots, \Sigma_4$, and the construction of Case II to Σ_5 . Composing the degenerations one by one, we obtain Figure 7 as our limit configuration. Now, since the limiting $\mathbb{P}^3$ ’s span a projective space of dimension $(4 \cdot 21 - 1)$, hence, by semicontinuity of the dimension we get a lower bound on $\dim(\text{Sec}_{20}(V_{3,6}))$, and we can conclude that $V_{3,6}$ is not 20–defective. $\square$

Lemma 4.2. $(\widetilde{\mathbb{P}^3})$ embedded in $\mathbb{P}^{80}$ via $\pi^*(\mathcal{O}_{\mathbb{P}^3}(8)) \otimes \mathcal{O}_{\widetilde{\mathbb{P}^3}}(-7E)$ is not 19–defective.




FIGURE 8. S_8^1

Proof. Figure 8 represents a layer S_8^1 , whose corresponding toric variety is $(\widetilde{\mathbb{P}^3})$ in its embedding in $\mathbb{P}^{80}$. In the polytope we have colored 20 tetrahedra of the type introduced in Lemma 2.10. Then the toric varieties they define are tangent $\mathbb{P}^3$ ’s. Since the tetrahedra are pairwise disjoint, the relative tangent $\mathbb{P}^3$ ’s are skew in the ambient space $\mathbb{P}^{80}$, so they span a linear space of maximal dimension $4 \cdot 20 - 1$. Notice that in the picture there is one point that is not included in any tetrahedron. If instead of the tangent $\mathbb{P}^3$ ’s defined by the tetrahedra in the picture we take general ones, by semicontinuity we still have that $\dim(\text{Sec}_{19}(\widetilde{\mathbb{P}^3}))$ is maximal.

□

Lemma 4.3. $(\widetilde{\mathbb{P}^3})$ embedded in $\mathbb{P}^{120}$ via $\pi^*(\mathcal{O}_{\mathbb{P}^3}(10)) \otimes \mathcal{O}_{\widetilde{\mathbb{P}^3}}(-9E)$ is not 29-defective.

Proof. In Figure 9 we see a layer S_{10}^1 which corresponds to the embedding of $(\widetilde{\mathbb{P}^3})$ in $\mathbb{P}^{120}$. The colored tetrahedra inside S_{10}^1 are 30, and only two of them are not of the type described in Lemma 2.10. Applying the argument of Lemma 4.1, we regard Figure 9 as the limit of a configuration composed of 30 independent tangent $\mathbb{P}^3$'s at general points of $(\widetilde{\mathbb{P}^3})$. The $\mathbb{P}^3$'s of the limiting configuration span a projective space of dimension $(4 \cdot 30 - 1)$. Then, again by semicontinuity of the dimension, we have that $\text{Sec}_{29}(\widetilde{\mathbb{P}^3})$ is not 29-defective. As well as in S_8^1 we have one point left out of the configuration. □





FIGURE 9. The layer S_{10}^1 : the two tetrahedra in blue do not correspond to tangent $\mathbb{P}^3$'s.

Corollary 4.4. $V_{3,8}$ and $V_{3,10}$ are not k -defective for any $k \in \mathbb{N}$.

Proof. Let us consider the configuration composed of the tetrahedron of Figure 7 leaning on a layer S_7^1 , which in turn lies on a layer S_8^1 as the one in Figure 8. We denote by D_8 such a configuration; it can be seen as the central fiber (composed of Δ_6 , S_7^1 and S_8^1) of a degeneration of $V_{3,8}$. By Lemma 4.1 and Lemma 4.2, we can apply Proposition 2.5, being Δ_6 and S_8^1 disjoint, thus we obtain that $\dim(\text{Sec}_{n_8}(V_{3,8})) = 4 \cdot 41 - 1 = 163$, i.e. $V_{3,8}$ is not 40-defective. Note that $\text{Sec}_{n_8}(V_{3,8}) \subset \mathbb{P}^{164} = \mathbb{P}^{N_8}$, hence $\text{Sec}_{n_8}(V_{3,8})$ is a hypersurface. We look now at the defectivity of $\text{Sec}_{n_8+1}(V_{3,8})$; let us denote by H_{n_8} the general tangent space to $\text{Sec}_{n_8}(V_{3,8})$, which is a hyperplane, and by H_{n_8+1} the tangent space to $\text{Sec}_{n_8+1}(V_{3,8})$ at a general point. Then, by Terracini's Lemma we get that $H_{n_8+1} = \langle H_{n_8}, T_p(V_{3,8}) \rangle$, p being a general point on $V_{3,8}$; hence $\dim(H_{n_8+1}) = N_8$. It follows that $V_{3,8}$ is not k -defective for any $k > n_8$, hence for any $k \in \mathbb{N}$ (see Lemma 2.7). We use a similar argument to show that $V_{3,10}$ is not n_{10} -defective: let us consider the configuration D_{10} composed of D_8 leaning on two layers, one of type S_9^1 , below which we put a S_{10}^1 as the one in Figure 9. Since D_8 and S_{10}^1 are disjoint polytopes, we apply again Proposition 2.5 to this configuration, using Lemma 4.3, thus we get that $V_{3,10}$ is not 70-defective. Now we want to extend the non k -defectivity of $V_{3,10}$ to any $k \in \mathbb{N}$. If $k < n_{10}$ we apply Lemma 2.7; let us now consider $\text{Sec}_{n_{10}+1}(V_{3,10})$, we call $H_{n_{10}}$ the general tangent space to $\text{Sec}_{n_{10}}(V_{3,10})$, and $H_{n_{10}+1}$ the tangent space to $\text{Sec}_{n_{10}+1}(V_{3,10})$ at some general point. Looking at the polytopes which compose D_{10} , we see that two vertices stay out of the configuration (one pointed out in D_8 , the other one in S_{10}^1); indeed $\dim(\text{Sec}_{n_{10}}(V_{3,10})) = N_{10} - 2$. Hence $\text{codim}(H_{n_{10}}) = 2$. Again by Terracini's lemma we get that $H_{n_{10}+1} = \langle H_{n_{10}}, T_p(V_{3,10}) \rangle$, with p a general point on $V_{3,10}$. Then $\dim(H_{n_{10}+1}) = N_{10}$; if it were not, then $H_{n_{10}+1}$ would be a hyperplane in $\mathbb{P}^{N_{10}}$; let $\pi_{n_{10}}$ be the projection from $H_{n_{10}}$ in $\mathbb{P}^{N_{10}}$ onto $\mathbb{P}^1$. Since $T_p(V_{3,10}) \subset H_{n_{10}+1}$, its projection is a point. But $\pi_{n_{10}}(T_p(V_{3,10}))$ is exactly the general tangent space to the projection of $V_{3,10}$, which must be a point. Hence $V_{3,10}$ would be degenerate, which is a contradiction. With this argument we get that $\text{Sec}_k(V_{3,10})$ is not k -defective for any $k > n_{10}$. □

Lemma 4.5. $V_{3,12}$ is not k -defective for any $k \in \mathbb{N}$.

FIGURE 10. The layers $S_9^1, S_{10}^1, S_{11}^1, S_{12}^1$ in D_{12}

Proof. Let us look at Figure 10; it is a composition of layers of type $S_9^1, S_{10}^1, S_{11}^1$ and S_{12}^1 . We place over the four layers represented, a configuration of type D_8 , as in Corollary 4.4. We call this new configuration D_{12} ; we are going to show that $V_{3,12}$ is not n_{12} -defective using the strategy of Lemma 4.1. Looking at the 113 colored tetrahedra in D_{12} (counting also the ones in $D_8 \subset D_{12}$), most of them correspond to tangent $\mathbb{P}^3$'s, as in Lemma 2.10, but 13 of them are not tangent $\mathbb{P}^3$'s, so we want to think of them as limits of general tangent $\mathbb{P}^3$'s. If we denote by $\Sigma_1, \dots, \Sigma_{13}$ the mentioned tetrahedra, we see that some of them lie inside unitary cubes, but we find three tetrahedra, one lying in D_6 ($D_6 \subset D_8$, moreover see Lemma 4.1), one in S_{10}^1 , the other one in S_{11}^1 , which sit inside blocks Σ as in Figure 3. Let us call them Σ_1, Σ_2 and Σ_3 . For all the tetrahedra $\Sigma_1, \dots, \Sigma_{13}$ we can reapply the techniques used in Lemma 4.1, i.e. one by one we compose the degenerations of general tangent $\mathbb{P}^3$'s to the limit $\mathbb{P}^3$'s defined by the relative Σ_i 's, $i = 1, \dots, 13$. Of course we will apply the construction of Case II to Σ_1, Σ_2 and Σ_3 , and the construction of Case I to $\Sigma_4, \dots, \Sigma_{13}$. At the end of the process, composing all the degenerations, we obtain D_{12} as our limit configuration. Finally we observe that, since the 113 limit $\mathbb{P}^3$'s span a projective space of dimension $(4 \cdot 113 - 1)$, hence, by semicontinuity of the dimension we get a lower bound on $\dim(\text{Sec}_{n_{12}}(V_{3,12}))$, and we can conclude that $V_{3,12}$ is not n_{12} -defective. But we are interested in the k -defectivity of $V_{3,12}$ for any $k \in \mathbb{N}$. So, we are going to study $\text{Sec}_{n_{12}+1}(V_{3,12})$. Let us notice that in the configuration D_{12} three points are

not contained in any of the pointed out tetrahedra. However they are very close to each other, so we can think of the three points as being contained in a further tetrahedron in D_{12} . Having in mind a configuration with $n_{12} + 2$ disjoint tetrahedra, we can repeat the argument we used before. Then again by semicontinuity we obtain that $\text{Sec}_{n_{12}+1}(V_{3,12})$ spans the whole ambient space. It follows that $V_{3,12}$ is not k -defective for any $k > n_{12}$, hence for any k (see Lemma 2.7). $\square$

So far we have shown that $V_{3,d}$ is not k -defective for $d = 6, 8, 10, 12$ and any $k \in \mathbb{N}$. In the following we will study the cases $d \equiv 6 \pmod{8}$, $d \equiv 0 \pmod{8}$, $d \equiv 10 \pmod{8}$, which we will handle using recurrence on the configurations we have already built.




FIGURE 11. Δ_{14}

4.1. Case $d = 6 + 8k$. We start by examining the case $d = 6 + 8k$. When $k = 1$, the tetrahedron Δ_{14} (see Lemma 2.9) can be split up into one Δ_6 , that we triangulate following Figure 7, and one layer S_{14}^7 of height 7 and big base of side length 14. We can furthermore divide S_{14}^7 in two triangulated tetrahedra Δ_6 lying in its ends, and a block Ξ , as illustrated in Figure 11. We can triangulate Ξ as follows: one semicube P_7 of side length 7, plus one tetrahedron of side length 5, that we denote by T_5 , and one tetrahedron of side length 6, T_6 . None of T_5 and T_6 is a 3-simplex, but we can think of them as the configurations of $V_{3,5}$ and $V_{3,6}$ respectively (the subdivision of Ξ is represented in Figure 12). Recall that $V_{3,5}$ and $V_{3,6}$ are respectively not 13-defective and 20-defective. In Figure 13(a) we have




FIGURE 12. Subdivision of the block Ξ

represented the orthogonal projection of the semicube P_7 , which shows that it contains 6 disjoint columns of height 7, each containing 4 pairwise disjoint unitary cubes, and every cube is not 1-defective; the squares in Figure 13(a) are precisely the projections of the columns. The yellow triangles correspond to the projections of some different columns of height 7, that we denote by γ_7 , each containing 6 pairwise disjoint tetrahedra, as in Figure 13(b). If we consider the toric variety defined by γ_7 , it is a $\mathbb{P}^1 \times \mathbb{P}^2$ embedded in $\mathbb{P}^{23}$ via $p_2^*(\mathcal{O}_{\mathbb{P}^2}(1)) \otimes p_1^*(\mathcal{O}_{\mathbb{P}^1}(7))$, where p_1, p_2 are the projections on the factors; we see that it is not 5-defective. To show this, we will use the techniques of Lemma 4.1. We want to prove that the 6 tetrahedra that we pointed out in γ_7 correspond to limits of general tangent $\mathbb{P}^3$'s. Four of them are not of the type introduced in Lemma 2.10; let us denote them by $\Sigma_1, \dots, \Sigma_4$; every Σ_i lies inside a semicube, which corresponds to $\mathbb{P}^1 \times \mathbb{P}^2$ and spans a $\mathbb{P}^5$. So in order to use the argument of Lemma 4.1 we just have to pay attention to the ambient space. An easy calculation such as in Lemma 4.1 yields that $\Sigma_1, \dots, \Sigma_4$ are limits of general tangent $\mathbb{P}^3$'s hence Figure 13(b) is our limit configuration. Notice that the tetrahedra corresponding to the limit $\mathbb{P}^3$'s are pairwise disjoint, then they span a projective space of dimension

$(4 \cdot 6 - 1)$. Hence, by semicontinuity we can conclude that the toric variety defined by γ_7 is not 5-defective. Going back to P_7 , since it is composed of 4 columns γ_7 plus $7 \cdot 4$ unitary cubes and all these polytopes are pairwise disjoint, applying Proposition 2.5 we get that the toric variety associated to P_7 is not 71-defective. Recalling that above S_{14}^7 lies a Δ_6 , which represents a $V_{3,6}$, and that in our configuration (see Figure 11) we have disjoint polytopes, then we can apply Proposition 2.5 and obtain that $V_{3,14}$ is not n_{14} -defective (where $n_{14} + 1 = 170$). In





FIGURE 13.

order to study the generic case $d = 6 + 8k$, we look at the tetrahedron Δ_{6+8k} ; we proceed by recurrence. Suppose we already know that $V_{3,d-8}$ is not n_{d-8} -defective. We build a configuration of Δ_d in the following way: we set a layer of height 7 with base of side length d , i.e. S_d^7 , below Δ_{d-8} , whose configuration we already know by induction; now we subdivide S_d^7 as follows: first we note that if $d = 6 + 8k$, then S_d^7 contains $\frac{k(k-1)}{2}$ cubes of side length 7 (C_7). In C_7 we insert 16 disjoint columns of height 7, each containing 4 disjoint unitary cubes, which, in turn, are not 1-defective, as shown in Lemma 2.11. Hence, being $16 \times 4 \times 2 = 128$, we get that C_7 is not 127-defective. Moreover, S_d^7 contains k semicubes equal to P_7 , that we triangulated before, and finally, on the skew stripe located in the front we set $k + 1$ copies of Δ_6 , k copies of T_5 and k ones of T_6 , as in Figure 14, which represents the orthogonal projection of S_{22}^7 . We summarize our construction in the following computation:

$$(4.2) \quad \Delta_d = \Delta_{d-8} + \left(\frac{k(k-1)}{2} \cdot C_7 \right) + (k \cdot P_7) + (2 \cdot \Delta_6) + k \cdot (T_5 + T_6) + (k-1) \cdot \Delta_6$$


FIGURE 14. Orthogonal projection of S_{22}^7 .

Hence, recalling the defectivity of the toric varieties associated to the polytopes composing Δ_d , we observe that

$$n_d = n_{d-8} + \frac{k(k-1)}{2} \cdot 128 + k \cdot 72 + 2 \cdot 21 + k \cdot (14 + 21) + (k-1) \cdot 21.$$

Now, noticing that the polytopes involved are all disjoint, we can apply Proposition 2.5, and we get that $V_{3,6+8k}$ is not n_{6+8k} -defective.

4.2. Configuration for $d = 8 + 8k$. The case $d = 8 + 8k$ with $k \geq 1$ is more complicated. As before, let $k = 1$. Then $d = 16$, and we can decompose Δ_{16} setting a Δ_8 above a S_{16}^7 -layer.

FIGURE 15. Subdivision of S_{16}^7 .

The problem is now to find a configuration for S_{16}^7 : we insert a prism having triangular basis of side length 9 and height 7, denoted P_9 ; it can be viewed as the half of a parallelepiped of height 7 having square basis of side 9, called C_9 ; then, on the skew stripe in the front we put two tetrahedra Δ_6 and a block B_8 composed of a T_6 plus a cut tetrahedron of side 7 which we denote T_7^* , as it is shown in Figure 15. We can obtain T_7^* setting a Δ_5 on a






FIGURE 16. The cut.

layer S_6^1 , in turn lying on a S_7^1 , from which we remove the black prism as shown in Figure 16(a). We want to show that T_7^* is not 27-defective. Notice that Δ_5 is not 13-defective; we focus on the cut S_7^1 , which is disjoint from Δ_5 . Let us consider the configuration in Figure 16(b). We see that 10 tetrahedra correspond to the ones introduced in Lemma 2.10, i.e. they are tangent $\mathbb{P}^3$'s. But there are 4 tetrahedra adjacent to the cut, which do not correspond to tangent $\mathbb{P}^3$'s. Each of them lies inside a semicube, so we apply the argument of Lemmas 4.1 and 4.5, that we also saw for the column γ_7 , and we get that the tetrahedra inside S_7^1 can be interpreted as limits of tangent $\mathbb{P}^3$'s. Being disjoint, they span a projective space of dimension $14 \cdot 4 - 1$, hence the cut layer S_7^1 is not 13-defective (again with abuse of notation). The block B_8 is not 48-defective. Summing up, by Proposition 2.5 we get that $V_{3,16}$ is not n_{16} -defective. In the general case, when $d = 8 + 8k$, we consider the parallelepiped C_9 , from which we remove a vertical edge of height 7: in this way we obtain a prism, H_9 , that we can see in Figure 17, and that contains 96 disjoint unitary cubes plus one column γ_7 . So H_9 is not 197-defective. Just like before we denote by P_7 the half

FIGURE 17. Orthogonal projection of H_9 .

of a cube C_7 ; P_7 is not 71-defective. We are now able to split Δ_d into Δ_{d-8} , and a S_d^7 . In this layer we insert (see Figure 18) in diagonal order: one H_9 ; 4 copies of P_7 ; then we have $\sum_{i=1}^{k-3} (i \cdot C_7 + 4 \cdot P_7)$; k copies of P_7 ; in the front we put $(k+1)$ copies of Δ_6 , $(k-1)$ copies of the block $T_5 + T_6$, and finally one block B_8 . Hence we have

FIGURE 18. Orthogonal projection of S_{40}^7 .

that:


$$\Delta_d = \Delta_{d-8} + H_9 + 4 \cdot P_7 + \sum_{i=1}^{k-3} (i \cdot C_7 + 4 \cdot P_7) + k \cdot P_7 + (k-1) \cdot (T_5 + T_6) + (k+1) \cdot \Delta_6 + B_8.$$

Substituting in the last expression the defectivity of each summand, we get:

$$n_d = n_{d-8} + 198 + 4 \cdot 72 + \sum_{i=1}^{k-3} (i \cdot 128 + 4 \cdot 72) + k \cdot 72 + (k-1) \cdot (14 + 21) + (k+1) \cdot 21 + 49.$$

So applying Proposition 2.5 we get that $V_{3,8+8k}$ is not n_{8+8k} -defective.

4.3. Configuration for $d = 10 + 8k$. Let us examine the case $k = 1$, corresponding to $d = 18$; the tetrahedron Δ_{18} is composed of a Δ_{10} leaning on a layer S_{18}^7 .

FIGURE 19. Subdivision of S_{18}^7 .

Let us subdivide S_{18}^7 (see Figure 19): we call P_{11} the half of a parallelepiped of height 7 and basis of side length 11; furthermore, in the front we set two copies of Δ_6 and a block B_{10} composed of a T_6 and a tetrahedron of side 9 from which we remove a prism, as we did for T_7^* . We obtain a solid, T_9^* ; we want to show that it is not 41-defective. Let us decompose T_9^* in the following way: a tetrahedron Δ_5 leaning on a layer S_6^1 , and this lies on a S_9^3 that we cut as in Figure 20(a). In particular we remove the part colored in black and blue, and in the remaining part we dispose the tetrahedra. Looking at Figure 20(a) we see that S_9^3 with its cut is composed of 3 layers, S_7^1 , S_8^1 and S_9^1 , all having a cut as well, so we dispose the tetrahedra in the S_7^1 and in S_9^1 as in Figure 16(b); indeed the blue

parts correspond to the black solid that we took off in Figures 16(a) and 16(b). Let us observe that the tetrahedra lying in T_9^* are 42. Repeating the argument used for T_7^* , it follows that the block B_{10} is not 62-defective. In P_{11}




FIGURE 20.

we insert 15 disjoint columns of height 7 and square basis of side 1, each containing 4 disjoint cubes; moreover P_{11} contains 6 columns γ_7 . Then P_{11} is not 155-defective. Summing up and using Proposition 2.5 we get that $V_{3,18}$ is not n_{18} -defective. In the general case $d = 10 + 8k$, as usual we divide Δ_d in one triangulated Δ_{d-8} plus one S_d^7 . In the layer S_d^7 we insert: $2(k-2)$ copies of C_7 , 2 copies of P_7 , one block B_{10} , $k+1$ copies of Δ_6 , $k-1$ copies of $T_5 + T_6$ and finally one block denoted by A_k . Let us show the configuration of this block: for $k = 2$, the block A_2 contains 33 disjoint columns of height 7 and square basis of side 1, and 2 columns γ_7 , hence it is not 275-defective. The orthogonal projections of these columns can be viewed in Figure 20(b). For $k \geq 3$ the block A_k is different: in Figure 20(c) we see the projection of A_3 . With this picture in mind, we observe that A_k contains columns of height 7 and side 1 (not 7-defective), and columns γ_7 (not 5-defective). In particular it contains $\left(35 + \sum_{i=3}^{4(k-2)+1} i\right)$ columns of the first type, and $(4 \cdot (k-2) + 2)$ columns γ_7 . Thus if we set: $\alpha_2 + 1 := 276$, and for $k \geq 3$, $\alpha_k + 1 := 8 \cdot \left(35 + \sum_{i=3}^{4(k-2)+1} i\right) + 6 \cdot [4 \cdot (k-2) + 2]$, we have that A_k is not α_k -defective for any $k \geq 2$. So we get that the decomposition in polytopes is the following (see Figure 21(a) and




FIGURE 21. Projections of subdivisions.

Figure 21(b)):

$$\Delta_d = \Delta_{d-8} + A_k + 2(k-2) \cdot C_7 + 2 \cdot P_7 + (k+1) \cdot \Delta_6 + B_{10} + (k-1) \cdot (T_5 + T_6).$$

The last expression corresponds to the following defectivities:

$$n_d = n_{d-8} + (\alpha_k + 1) + 2(k-2) \cdot 128 + 2 \cdot 72 + (k+1) \cdot 21 + 63 + (k-1) \cdot (14 + 21).$$

Hence we conclude that $V_{3,10+8k}$ is not n_{10+8k} -defective.

4.4. Configuration for $d = 12 + 8k$. This is the last configuration, and it is similar to the one proposed for $d = 10 + 8k$; again we start examining the case $k = 1$ corresponding to $d = 20$. We decompose Δ_{20} into one tetrahedron Δ_{12} and one S_{20}^7 . The subdivision of S_{20}^7 is the following: one prism of height 7 with triangular basis of cathetus 13, that we call P_{13} ; in the front lie 2 copies of Δ_6 , one block B_{12} composed of a T_6 and a solid T_{11}^* obtained removing a piece from Δ_{11} as in Figure 22. As we already did for T_7^* and T_9^* , we can think of T_{11}^* as




FIGURE 22. Subdivision of S_{20}^7 .

obtained by cutting a Δ_{11} . So we get a solid composed of a Δ_5 leaning on a layer S_6^1 in turn leaning on a S_{11}^5 that has been cut as in Figure 23(a), where we remove the part in black and blue. Using again the triangulation shown




FIGURE 23.

in Figure 16(b) behind the removed blue parts, we see that T_{11}^* is not 55-defective, and this implies that the block B_{12} is not 76-defective (where $21 + (91 - 35) = 77$). The decomposition in polytopes is the following:

$$\Delta_{20} = \Delta_{12} + P_{13} + 2 \cdot \Delta_6 + B_{12},$$

that is to say

$$n_{20} = n_{12} + 210 + 2 \cdot 21 + 77 = 442.$$

Then $V_{3,20}$ is not n_{20} -defective. In order to discuss the general case $d = 12 + 8k$, we consider the subdivision of Δ_d in a Δ_{d-8} plus a S_d^7 . Then we insert in S_d^7 a block A_k , whose configuration we will discuss in the following. For $k = 2$, the block A_2 contains 43 disjoint columns of height 7 and square basis of side 1, and 3 columns γ_7 , hence it is not 361-defective. The orthogonal projections of these columns can be viewed in Figure 23(b). For $k \geq 3$ the block A_k is different: in Figure 23(c) we see the projection of A_3 . Looking at this picture, again we see that A_k contains columns of height 7 and side 1 (not 7-defective), and columns γ_7 (not 5-defective). In particular it contains $\left(46 + \sum_{i=4}^{4k-6} i\right)$ columns of the first type, and $(4k - 5)$ columns γ_7 . As well as in the previous case we set: $\alpha_2 + 1 := 362$, and for $k \geq 3$, $\alpha_k + 1 := 8 \cdot \left(46 + \sum_{i=4}^{4k-6} i\right) + 6 \cdot (4k - 5)$, hence we have that A_k is not

α_k -defective for any $k \geq 2$. In addition to A_k , in S_d^7 we insert: $2(k-2)$ copies of C_7 , 2 copies of P_7 , and in the front stripe there are a block B_{12} , $(k+1)$ copies of Δ_6 plus $(k-1)$ copies of $(T_5 + T_6)$. Summarizing, we have that:

$$\Delta_d = \Delta_{d-8} + A_k + 2(k-2) \cdot C_7 + 2 \cdot P_7 + (k+1) \cdot \Delta_6 + B_{12} + (k-1) \cdot (T_5 + T_6).$$

One example of this subdivision is available for S_{36}^7 , whose projection can be viewed in Figure 24. Substituting in the last expression the defectivities, we get:

$$n_d = n_{d-8} + (\alpha_k + 1) + 2(k-2) \cdot 128 + 2 \cdot 72 + (k+1) \cdot 21 + 77 + (k-1) \cdot (14 + 21).$$




FIGURE 24. Orthogonal projection of S_{36}^7 .

Hence we have that $V_{3,12+8k}$ is not n_{12+8k} -defective. This completes the proof of Alexander-Hirschowitz's theorem in dimension 3.

Remark 4.6. Let us observe that in fact we have proved something more: we have decomposed the polytopes Δ_d corresponding to $V_{3,d}$, into smaller subpolytopes, and in order to prove the not-defectivity of $V_{3,d}$, we have shown that some of those subpolytopes correspond to not-defective toric varieties; this is further information due to this approach.

REFERENCES

- [BO07] M. C. Brambilla, G. Ottaviani, On the Alexander-Hirschowitz Theorem, *Journal of Pure and Applied Algebra*, Vol. 212 (2008), no. 5, 1229-1251
- [Ch02] K. A. Chandler, A brief proof of a maximal rank theorem for generic double points in projective space, *Trans. Amer. Math. Soc.* 353 (2001), no. 5, 1907-1920
- [Ci01] C. Ciliberto, Geometric Aspects of Polynomial Interpolation in More Variables and of Waring's Problem, *Progress in Mathematics*, Vol 201, 2001 Birkhäuser Verlag Basel/Switzerland
- [CM06] C. Ciliberto, O. Dumitrescu e R. Miranda, Degenerations of the Veronese and Applications, 2006, preprint
- [DC05] D. A. Cox, LECTURES ON TORIC VARIETIES, 2005, indirizzo URL: <http://www.cs.amherst.edu/~dac/lectures/coxcimpa.pdf>
- [JD06] Jan Draisma, A tropical approach to secant dimensions, *Mathematics*, abstract math.AG/0605345, 2006
- [EH00] D. Eisenbud, J. Harris, *The Geometry of Schemes*, Springer-Verlag, New York, 2000
- [Fu93] W. Fulton, *Introduction to Toric Varieties*, Princeton University Press, Princeton, New Jersey, 1993
- [GR90] R. C. Gunning, *Introduction to holomorphic functions of several variables*, Wadsworth & Cole Mathematics Series, Belmont, California, 1990
- [Ha97] R. Hartshorne, *Algebraic Geometry*, Springer-Verlag, New York Berlin Heidelberg, 1997
- [Hu01] S. Hu, Semi-Stable Degenerations of Toric Varieties and Their Hypersurfaces, *Mathematics*, abstract math.AG/0110091 v1 8 Oct 2001, University of Wisconsin-Madison, 2001
- [KY91] K. Yang, *Complex Algebraic Geometry, An Introduction to Curves and Surfaces*, MARCEL DEKKER, INC., New York, 1991
- [Mi04] R. Miranda, *Anacapi Lectures On Degenerations Of Surfaces*, Anacapi, 2004

- [Od88] T. Oda, Convex Bodies and Algebraic Geometry—An introduction to the Theory of Toric Varieties, Springer–Verlag, Berlin Heidelberg, 1988
- [Sh77] I. Shafarevich, Basic Algebraic Geometry, Springer–Verlag, Berlin Heidelberg New York, 1977
- [SS05] B. Sturmfels, S. Sullivant, Combinatorial secant varieties, [math.AC/0506223](https://arxiv.org/abs/math.AC/0506223)

S. BRANNETTI: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA 3, LARGO SAN LEONARDO MURIALDO, 1, 00146 ROMA, ITALY

E-mail address: `brannett@mat.uniroma3.it`